

Upneeq™

(oxymetazoline hydrochloride
ophthalmic solution) 0.1%

UPNEEQ has arrived!



Not an actual patient.

ONE DROP DAILY. UPLIFTING RESULTS.

**UPNEEQ is the first and only FDA-approved
eyedrop for treatment of droopy eyelid
(acquired blepharoptosis) in adults**

Learn more about **UPNEEQ**—the drop that lifts

IMPORTANT SAFETY INFORMATION

INDICATION

UPNEEQ™ (oxymetazoline hydrochloride ophthalmic solution), 0.1% is indicated for the treatment of acquired blepharoptosis in adults.

WARNINGS AND PRECAUTIONS

- Alpha-adrenergic agonists as a class may impact blood pressure. Advise UPNEEQ patients with cardiovascular disease, orthostatic hypotension, and/or uncontrolled hypertension or hypotension to seek medical care if their condition worsens.

Please see Important Safety Information continued on next page.

Please see accompanying Full Prescribing Information and Patient Information.

Ask Your Eye Doctor About

Upneeq™

(oxymetazoline hydrochloride
ophthalmic solution) 0.1%



A whole new way to treat
droopy eyelid (acquired
blepharoptosis)

Not an actual patient.

What can UPNEEQ do?

- ***It's the drop that lifts***—a unique daily medication designed to lift droopy eyelids
- ***It can improve field of vision***—which may help you see better
- ***It can lift your eyelid(s) quickly***
- ***Use UPNEEQ once a day, every day*** to improve field of vision and elevate your eyelids

IMPORTANT SAFETY INFORMATION (continued)

WARNINGS AND PRECAUTIONS

- Use UPNEEQ with caution in patients with cerebral or coronary insufficiency or Sjögren's syndrome and advise patients to seek medical care if signs and symptoms of potentiation of vascular insufficiency develop.
- UPNEEQ may increase the risk of angle closure glaucoma in patients with untreated narrow-angle glaucoma. Advise patients to seek immediate medical care if signs and symptoms of acute narrow-angle glaucoma develop.
- Patients should not touch the tip of the single patient-use container to their eye or to any surface, in order to avoid eye injury or contamination of the solution.

ADVERSE REACTIONS

- Adverse reactions that occurred in 1-5% of subjects treated with UPNEEQ were punctate keratitis, conjunctival hyperemia, dry eye, blurred vision, instillation site pain, eye irritation and headache.

Please see important safety information continued on back cover.

How can UPNEEQ benefit me?

Using UPNEEQ every day can provide an improved field of vision, which may help you see better, along with a visible lift in your eyelid's position.



Not an actual patient.

How often should I use UPNEEQ?

You may think that once your eyelid's position is elevated, you don't need to use UPNEEQ anymore. But UPNEEQ is not a "cure." For continued benefits, you need to use it once a day, every day.

Please see accompanying Full Prescribing Information and Patient Information.


Upneeq™

(oxymetazoline hydrochloride
ophthalmic solution) 0.1%

Apply UPNEEQ in 2 easy steps



Cut open foil wrapper and remove single-use vial



Apply UPNEEQ drops as directed, once a day



Don't touch your eye with the tip of the vial



Throw the vial away after applying drop(s)

If you wear contact lenses, remove them before applying UPNEEQ eyedrops. You may put them back in 15 minutes after applying UPNEEQ.

If more than one topical ophthalmic drug is being used, the drugs should be administered at least 15 minutes between applications.

For complete instructions on using UPNEEQ, please see the Instructions for Use section in the full Prescribing Information.

Visit [MeetUpneeq.com](https://www.meetupneeq.com) and learn more.

**UPNEEQ is available exclusively through the RVL Pharmacy
844-RVL-EYES (844-785-3937)**

DRUG INTERACTIONS

- Alpha-adrenergic agonists, as a class, may impact blood pressure. Caution in using drugs such as beta blockers, anti-hypertensives, and/or cardiac glycosides is advised. Caution should also be exercised in patients receiving alpha adrenergic receptor antagonists such as in the treatment of cardiovascular disease, or benign prostatic hypertrophy.
- Caution is advised in patients taking monoamine oxidase inhibitors which can affect the metabolism and uptake of circulating amines.

To report SUSPECTED ADVERSE REACTIONS or product complaints, contact RVL Pharmaceuticals at 1-877-482-3788. You may also report SUSPECTED ADVERSE REACTIONS to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see accompanying Full Prescribing Information and Patient Information.

Distributed by: RVL Pharmaceuticals, Inc.
Bridgewater, NJ 08807
Customer Service 1-866-600-4799

RVL
PHARMACEUTICALS, INC.

©2020 RVL Pharmaceuticals, Inc.
PM-US-UPN-0009